

2020-06-10

## Estimates of Sodium and Potassium Intake and Their Molar Ratio in Irish Children: Preliminary Findings From NCFSII

Eoin James Morrissey

*Department of Biological Sciences, Cork Institute of Technology, Cork, Ireland, eoin.morrissey@mycit.ie*

Laura Kehoe

*University College Cork, Cork, Ireland*

Aoibhin Moore Heslin

*University College Dublin, Dublin, Ireland*

Maria Buffini

*University College Dublin, Dublin, Ireland*

Breige McNulty

*University College Dublin, Dublin, Ireland*

*See next page for additional authors*

Follow this and additional works at: <https://sword.cit.ie/dptbiosciart>



Part of the [Biochemistry, Biophysics, and Structural Biology Commons](#), and the [Bioinformatics Commons](#)

---

### Recommended Citation

Morrissey, E., Kehoe, L., Heslin, A., Buffini, M., McNulty, B., Nugent, A., . . . Walton, J. (2020). Estimates of Sodium and Potassium intake and their molar ratio in Irish children: Preliminary findings from NCFSII. *Proceedings of the Nutrition Society*, 79(OCE2), E162. doi:10.1017/S002966512000110X

This Article is brought to you for free and open access by the Biological Sciences at SWORD - South West Open Research Deposit. It has been accepted for inclusion in Department of Biological Sciences Publications by an authorized administrator of SWORD - South West Open Research Deposit. For more information, please contact [sword@cit.ie](mailto:sword@cit.ie).

---

**Authors**

Eoin James Morrissey, Laura Kehoe, Aoibhin Moore Heslin, Maria Buffini, Breige McNulty, Anne Nugent, John Kearney, Albert Flynn, and Janette Walton

The 13th European Nutrition Conference, FENS 2019, was held at the Dublin Convention Centre, 15–18 October 2019

## Estimates of Sodium and Potassium intake and their molar ratio in Irish children: Preliminary findings from NCFSES

Eoin Morrissey<sup>1</sup>, Laura Kehoe<sup>2</sup>, Aoibhin Moore Heslin<sup>3</sup>, Maria Buffini<sup>3</sup>, Breige McNulty<sup>3</sup>, Anne Nugent<sup>4</sup>, John Kearney<sup>5</sup>, Albert Flynn<sup>2</sup> and Janette Walton<sup>1</sup>

<sup>1</sup>Cork Institute of Technology, Cork, Ireland,

<sup>2</sup>University College Cork, Cork, Ireland,

<sup>3</sup>University College Dublin, Dublin, Ireland,

<sup>4</sup>Queens University Belfast, Belfast, United Kingdom and

<sup>5</sup>Technological University Dublin, Dublin, Ireland

### Abstract

The Food Safety Authority of Ireland (FSAI) have set target maximum daily salt intakes for children (4–6y: 3 g, 7–10y: 5 g, 11–14y: 6g) while the European Food Safety Authority (EFSA) have set Adequate Intakes (AI) for potassium of 1100mg/d, 1800mg/d and 2700mg/d for children of the same respective age groups. An individual's sodium to potassium (Na:K) intake ratio is an important predictor of hypertension and the World Health Organization (WHO) recommend a Na:K intake ratio of  $\leq 1.0$ mmol/mmol for both adults and children. Although the morbidities associated with hypertension may not be seen until adulthood, blood pressure in childhood has a significant association with blood pressure in adulthood. Therefore, estimation of Na:K intake ratios (best measured by urinary excretion) in children may predict their susceptibility to hypertension related diseases in later life. The aim of this study was to estimate sodium and potassium intake and mean molar Na:K intake ratio of Irish children and to assess compliance with dietary guidance.

Morning spot urine samples were collected for 572 children aged 5–12 years (95% of total sample) as part of the nationally representative Irish National Children's Food Survey II (2017–2018) (NCFSES; www.iuna.net). Samples were transported, processed and stored using best practice procedures. Urinary excretion of sodium and potassium were measured using a Randox RX Daytona and were corrected for gender and age-specific 24-hour urine volume estimations based on 24-hour urine volume estimates from Australian children. SPSS Version 25 was used for all analyses.

Mean 24-hour urinary sodium excretion was 2018mg/d, equivalent to an average salt excretion of 5.0g/d exceeding the FSAI maximum target intake for all age groups except 11–12 year olds. Mean 24-hour urinary potassium excretion was 1411mg/d with mean intakes below the AI from EFSA for all age groups with the exception of 5–6 year olds. The mean molar Na:K ratio of Irish children was 2.8 for boys and 3.4 for girls. Only 5% of Irish children met the WHO recommendation for a Na:K ratio of  $\leq 1.0$ mmol/mmol.

High intakes of sodium and low intakes of potassium reported in this study result in a low compliance with the WHO recommendation of a Na:K ratio  $\leq 1.0$ mmol/mmol. This may lead to a higher risk of hypertension and related morbidities in later life. Based on these findings, dietary interventions to combat hypertension related diseases (such as lowering sodium and increasing potassium intakes) should be implemented from childhood.

### Conflict of Interest

There is no conflict of interest